

# Early reporting of efficacy endpoints and its potential impact: Clinical part

F. Cardoso, MD

Director, Breast Unit, Champalimaud Clinical Center, Lisbon, Portugal ESMO Board of Directors & NR Committee Chair ESO Breast Cancer Program Coordinator EORTC Breast Group Chair













#### **DISCLOSURES**

### Consultant/Ad Board:

Astellas/Medivation, AstraZeneca, Celgene, Daiichi-Sankyo, Eisai, GE Oncology, Genentech, GlaxoSmithKline, Macrogenics, Merck-Sharp, Merus BV, Novartis, Pfizer, Pierre-Fabre, Roche, Sanofi, Teva



## RISKS OF EARLY STOPPING/REPORTING TRIALS

- High probability of overestimation of the magnitude of treatment effect
- Moderate probability of underestimation of the magnitude of treatment effect (long natural history; e.g. RT effects in breast cancer)
- Repeated interim analyses at short intervals raise concern about data reliability
- Long-term benefits and long term toxicities unknown/partially known
- Lack of safety data if enrollment is stopped
- Impact of crossover on overall survival results, if allowed
- Influence on other ongoing trials



# Stopping a trial early in oncology: for patients or for industry?

F. Trotta<sup>1</sup>, G. Apolone<sup>2</sup>, S. Garattini<sup>2</sup> & G. Tafuri<sup>1,3</sup>\*

<sup>1</sup>Italian Medicines Agency (AIFA), Rome; <sup>2</sup>Mario Negri Institute for Pharmacological Research, Milan, Italy; <sup>3</sup>Utrecht University, Utrecht Institute for Pharmaceutical Sciences, Utrecht, The Netherlands

# Between 1997-2007, 25 RCTs testing new anti-cancer drugs stopped early because of benefit at interim analysis

- Result of early report: stop enrollment 48%; disclosure of results 20%; crossover to treatment group 12%; crossover to treatment group+ stop enrolment: 12%
- Increase in > 50% of prematurely stopped trials from 2004 to 2007
- 3300 patients/events across all studies were spared
- More than 78% of the RCTs were registration trials



STATISTICS IN ONCOLOGY

Stopping or Reporting Early for Positive Results in Randomized Clinical Trials: The National Cancer Institute Cooperative Group Experience From 1990 to 2005


Edward L. Korn, Boris Freidlin, and Margaret Mooney

From 1990-2005, 27 studies from National Cancer Institute Cooperative Group Trials stopped early due to a positive result.


In 17 trials, the treatment effect at last FU was similar or slightly smaller than the one reported earlier.

Trial Identifier	Treatment Effect	95% CI for HR or Difference in Rates	P
CLB-9011	Median OS: 66 v 56 months	Not applicable	.10
SWOG-8892	5-year OS: 67% v 37%	Not available	.001
E2491	5-year OS: 69% v 45%; difference = 24%	13.4% to 34.6%	.0001
SWOG-8814	HR = 0.83	0.69 to 0.99	.04
CLB-9344	HR = 0.82	0.71 to 0.95	.0064
CCG-5942	3-year OS: 98% $v$ 99%; difference = $-1$ %	-3.3% to 1.3%	.90
RTOG-9413	4-year OS: 84.7% v 84.3%; difference = 0.4%	-4.6% to 5.4%	<mark>.94</mark>
SWOG-S9701	HR = 0.84	0.61 to 1.16	.30
NCCTG-9741	HR = 0.66	0.54 to 0.82	.0001
NCIC-MA17	HR = 0.82	0.57 to 1.19	.3
ECOG-1496	HR = 0.51	0.25 to 1.04	.06
CCG-1961	HR = 0.64	0.47 to 0.87	.005
E2997	2-year OS: 79% $v$ 80%; difference = -1%	-14.3 to 12.3*	<mark>.69</mark>
NSABP-B-31/NCCTG-N9831	HR = 0.63	0.49 to 0.81	.0004
ECOG-2100	HR = 0.88	0.74 to 1.05	<mark>.16</mark>
NCIC-MA21	47 v 65 deaths	Not applicable	.09†


#### **MOLECULAR (RE-)CLASSIFICATION OF BREAST CANCER**



Perou et al, Nature, 2000 Sorlie et al, PNAS 2001 Sortiriou, PNAS 2003 Hu et al, BMC, Genomics 2006 Herschkowitz et al, GB 2007 Parker et al, JCO 2009





## **Annual Risk of Recurrence by ER Status**



- Over half of breast cancer recurrences occur >5 years post-surgery!
- The annual risk of late recurrence is particularly high in ER+ tumors (5.2% between years 5 and 8, 4.6% between years 8 and 12).


# More than Half of all Breast Cancer Recurrences and Deaths Occur Post- 5y Tamoxifen







## MA.17: Trial Design



- Eligibility criteria: postmenopausal at randomization, HR+, recurrence-free, completed 4.5–6 year Tamoxifen, ECOG PS 0–2
- Primary endpoint: DFS (breast- only events)
- Secondary endpoints: OS, rate of contralateral BC, safety, QoL,
- Sub-studies: BMD/bone markers, lipid profile



## **MA.17: Reports**

Year	Author	Journal	Title
2003	Goss et al	NEJM	A randomised trial of letrozole in postmenopausal women after five years of tamoxifen therapy for early- stage breast cancer
2005	Goss et al	JNCI	Randomized trial of letrozole following tamoxifen as extended adjuvant therapy in receptor-positive breast cancer: Updated findings from NCIC CTG MA.17
2008	Ingle et al	AnnOnc	Intent-to-treat analysis of the placebo-controlled trial of letrozole for extended adjuvant therapy in early breast cancer: NCIC CTG MA.17
2008	Goss et al	JCO	Late extended adjuvant treatment with letrozole improves outcome in women with early-stage breast cancer who complete 5 years of tamoxifen
2012	Jin et al	JCO	Longer-Term Outcomes of Letrozole Versus Placebo After 5 Years of Tamoxifen in the NCIC CTG MA.17. Trial: Analyses Adjusting for Treatment Crossover





### 2003: first results

# Median follow-up 2.4 y 207 events (40%)

Event	HR	95%CI	p value
DFS 4y	0.57	0.43- 0.75	.00008
OS 4y	0.61	0.47-0.79	< .001

# **Unblinding Expeditious report**







### Planned Subgroup analysis according to LN status

	DFS	Distant DFS	OS
Node + patients	HR 0.61*	HR 0.53*	HR 0.61*
	(95% CI, 0.45-0.84)	(95% CI, 0.36-0.78)	(95% CI, 0.38-0.98)
Node –	HR 0.45*	HR 0.63	HR 1.52
patients	(95% CI, 0.27-0.75)	(95% CI, 0.31-1.27)	(95% CI, 0.76-3.06)

> A similar reduction in local recurrences, new primaries, and distant recurrences occurred in N – and N+ patients, with OS benefit observed only in N-positive patients.

Extremely low rate rate Not surprising! than 70 years.

They relapse earlier; therefore: higher rate of events



### **Consequences of MA-17 early reporting**

**Trial stopped** 

**Patients offered crossover** 

Similar trials closed before completing accrual

Fortunately, FU of patients continued



### 2005 Update: 247 events (only 40 more events!)

### **Letrozol benefit for DFS 4y**


94.4% versus 89.8%


HR= 0.58 (95%CI: 0.45-0.76) (similar to 0.57)

**Absolute reduction in recurrence 4.6%** 

Treatment with extended Al received Guideline recommendation

#### No OS difference





Favors Letrozole

Favors Placebo



### 2008 Update: Intent to treat analysis


#### Median follow-up 64 months (over 5 years)

#### **DFS 4y benefit**

94.3% versus 91.4%: 2.9% absolute benefit (LOWER)

HR= 0.68 (95%CI: 0.55-0.83) (LOWER than before 0.58)

### No OS difference (95.1% in both)





# **2008 Update:** Cohort analysis of patients on Placebo who were offered Letrozol

**2012 Update: Long-term outcomes: <u>exploratory</u> analysis adjusting treatment crossover** 

#### Median FUP 64 months and 60% of crossover

Inverse probability of censoring weighted (IPCW) Cox model/ Cox model with time-dependent covariates

DFS HR= 0.52 (95%CI: 0.45-0.61)

OS HR= 0.61 (95%CI: 0.52-0.71)

No definite conclusion about OS



### **Consequences of MA-17 early reporting**

## B-33: Accrual

Opened: May, 1 2001

Target Accrual: 3000 pts

Accrual in 10/03: 1598 pts (53.3%)

Accrual stopped in October 2003 after disclosure of results from the NCIC MA.17 trial



### **Consequences of MA-17 early reporting**

# B-33 Plan After Disclosure of NCIC MA-17 Results


- In October 2003 the NSABP DMC decided to permanently stop accrual and to unblind the study
- Offer exemestane to patients in the placebo group free of charge
- Offer continuation of exemestane to patients in the exemestane group for a total of 5 years
- Notification of investigators and patients
- Rapid amendment of the study to implement the DMC recommendations



# **B-33:Trial Status Post Unblinding**

- Accrual of 3,000 pts was needed to detect a 21.3% reduction in hazard rate with a power of 80% (two-sided 0.05-level log-rank test).
- At the time of unblinding, 1598 pts had been randomized (1577 eligible)
- Upon unblinding, 560 of 783 pts on EXE continued EXE (72%)
- Of 779 pts on PLAC, 344 switched to EXE (44%)
- Median follow up for this ITT analysis: 30 mos

### **Consequences of MA-17 early reporting**





### **Consequences of MA-17 early reporting**


ATTENTION: Stopping the trial was a IDMC recommendation and was as per definition in the protocol

**WOULD YOU HAVE STOPPED THE TRIAL?** 



# Consequences of early reporting in CHEMOPREVENTION






Very large study, with potential to provide definite conclusions BUT reported earlier and stopped



# Consequences of early reporting in CHEMOPREVENTION




**CONSEQUENCIES:** 

- 1) The effect of chemoprevention on the incidence of breast cancer is known but not the true effect on mortality
- 2) Overall uptake of chemoprevention is low



# Consequences of early reporting <u>& subgroup</u> analysis in TRANSLATIONAL RESEARCH



3<sup>rd</sup> EXAMPLE



- NOT CONFIRMED IN BIG
   1-98 nor TEAM STUDIES
- Never reproduced

#### **CONSEQUENCIES:**

It influenced reimbursement and availability of Als in Belgium for years




#### IS IT ALWAYS WRONG TO STOP A TRIAL EARLIER?

What happens when the new treatment benefit is substantially superior?



# NCCTG N9831/NSABP B-31 trials: effect of adjuvant trastuzumab






### NCCTG N9831/NSABP B-31 trial

- Trial starting date May 2003 vs End of enrolment Nov 2004 vs Date of publication Oct 2005
- 81% of Patients accrued When Trial Stopped/Reported
- DSMC yes
- Sample size: 710 Planned vs 394 at Interim Analysis
- Primary endpoint used in interim analysis: DFS
- Interim analysis consequences: Stop enrolment and disclosures of results




### NCCTG N9831/NSABP B-31 trial: OS



Romond et al. N Engl J Med 2005;353:1673-84.

% Information	Treatment effect	P
55	HR=0,67	.015



Perez et al. J Clin Oncol 32:3744-3752.

% Information	Treatment effect	P
99	HR=0,63	<.001

Updated analysis confirmed the results.

Early reporting allowed for fewer patients to be treated in the control arm, crossover, earlier approval of drug.



# EARLY REPORTING OF TRIALS RISKS & BENEFITS



- 1) Expose less patients to the least effective therapy
- 2) Expose less patients to unnecessary toxicity
- Offer patients assigned to the control arm the better therapy, through crossover
- 4) Save resources that can be redirected to other relevant questions



- 1) Probability of overestimate the true treatment effect
- 2) Probability to underestimate the true treatment effect
- B) Loss of possibility to assess accurately long term outcomes such as OS and late toxicities
- 4) Implications for other ongoing trials

### **VERY BIG THANK YOU!**



Joana Ribeiro, MD



Berta Sousa, MD